

---


MG15 - Fifteenth Marcel Grossmann Meeting

# DECIGO : Gravitational-Wave Observation from Space

**Masaki Ando** (Univ. of Tokyo / NAOJ)  
on behalf of the DECIGO collaboration

# Outline


- Introduction
- B-DECIGO
- DECIGO





# First Detection of GW


- On Feb. 11<sup>th</sup>, 2016, **LIGO** announced **first detection of gravitational wave**. The signal was from inspiral and merger of **binary black hole**.
  - ⇒ Opens a new field of '**GW astronomy**'.



Courtesy Caltech/MIT/LIGO Laboratory

# Mergers of Binary Black Hole

- 2nd: GW151226 (2016.6 announce)
  - 3rd: GW170104 (2017.6.2 announce)
  - 4th: GW170814 (2017.9.27 announce)
  - 5th: GW170608 (2017.11.15 announce)
- Mergers of binary black holes would be **common events** in the universe.




<http://ligo.org/detections/GW170608>

LIGO/VIRGO



# Merger of Binary Neutron Stars

- On **Oct. 16<sup>th</sup>, 2017**, LIGO-VIRGO collaboration announced the first detection of gravitational-wave signal from merger of binary neutron stars
- The signal was detected on August 17<sup>th</sup>, 2017.  
→ Named **GW170817**.
- Source Localization  **$\sim 30\text{deg}^2$**




Courtesy Caltech/MIT/LIGO Laboratory

- **EM counterpart** was observed for the first time in GW170817.



- New knowledge
  - \* Origin of **SGRB**.
  - \* Origin of **heavy elements** in the universe.
  - \* **EoS** of neutron star
  - \* **Fundamental physics and cosmology**: speed of GW, Hubble's constant, ...

ApJL 848 L12 (2017)





# After the First Detections ...

---


- The first GW (and EM counter part) detections demonstrated new possibilities by **GW astronomy**, and also showed new mysteries, such as the origin of heavier mass ( $30M_{\odot}$ ) BBH.



- Network of **2<sup>nd</sup>-gen. GW antennae** (aLIGO, AdVIRGO, KAGRA, LIGO-India) will be formed in several years.
- Two ways after that for Astronomy and Cosmology:
  - **3<sup>rd</sup>-gen. ground-based GW antennae** (ET, CE).
  - **Space GW antennae** (LISA, DECIGO, TianQin, ...).


# Updated Roadmap for DECIGO

Figure: S.Kawamura





# B-DECIGO



# Space GW Observatory: B-DECIGO

※ We changed the name: Pre-DECIGO → B-DECIGO

- B-DECIGO

- Space-borne GW antenna formed by three S/C
- Target Sensitivity for GW :  $2 \times 10^{-23} \text{ Hz}^{-1/2}$  at 0.1Hz.

- Sciences of B-DECIGO

- (1) Compact binaries.
- (2) IMBH merger.
- (3) Info. of foregrounds for DECIGO.



Fig. by S.Sato

Target: JAXA Strategic Medium-scale mission (2020s).




# B-DECIGO Design (Preliminary)

- Mission Requirement

- Strain sensitivity of  $2 \times 10^{-23} \text{ Hz}^{-1/2}$  at 0.1Hz.
- >3-years observation period.

- Conceptual Design

- Laser interferometer by 3 S/C
- Baseline : 100 km
- Laser source : 1W, 515nm
- Mirror : 300mm, 30kg
- Drag-free and Formation flight.
- Record-disk orbit around the earth:  
Altitude 2000km, Period  $\sim 120\text{min}$  (Preliminary).




# Space GW antenna

## LISA

(Laser Interferometer  
Space Antenna)


- Target: SMBH, Binaries.  
GWs around 1mHz.
- Baseline : 2.5M km.  
Constellation flight by 3 S/C
- Optical transponder.



## B-DECIGO

(Deci-hertz Interferometer  
Gravitational Wave Observatory)

- Target: IMBH, BBH, BNS.  
GWs around 0.1Hz.
- Baseline : 100 km.  
Formation flight by 3 S/C.
- Fabry-Perot interferometer.






# Interferometer Configurations

- Optical transponder

- \* Phase locking of laser sources in each S/C.
- \* Long baseline is possible.
  - Better Acc. Noise

Doppler tracking  
using laser beam



M Hewitson (ASTROD WS 2017)

- Fabry-Perot cavity

- \* Direct reflection.
- \* Large laser power accumulated in cavity.
  - Better Shot noise

Similar config.to ground-based GW antennae





Fig: S. Kawamura

# Sensitivity Curves

T. Nakamura et al., Prog. Theor. Exp. Phys. 093E01 (2016)





# Sciences by B-DECIGO


---

- (1) Inspiral of Compact binaries [‘Promised’ target]
  - High rate  $\sim 10^5$  binaries/yr.
  - Estimation of binary parameters and merger time.
  - Astronomy by GW only and GW-EM observations.
- (2) Inspirals and mergers of IMBHs [Original science]
  - Cover most of the universe.
  - Formation history of SMBH and galaxies.
- (3) Foreground understandings for DECIGO [Cosmology]
  - Parameter estimation and subtraction of binaries.
  - Characteristics of foreground.
  - Is there any eccentric binaries?

# Target (1) : Compact Binaries

B-DECIGO will observe  $>100/\text{yr}$  binary NS inspirals.

$\sim 10^5/\text{yr}$  binary BH inspirals.




Low.-freq.  $\rightarrow$  **B-DECIGO**  
Mass, Position, Time,...

High-freq.  $\rightarrow$  Ground based  
Astrophysics, EoS of NS



# Sensitivity Curves

T. Nakamura et al., Prog. Theor. Exp. Phys. 093E01 (2016)



# Target (2) : Intermediate-mass BH Merger

B-DECIGO will see almost the whole Universe.



The mystery on the history of SMBH at the centers of Galaxies:


- (A) Large BH + Accretion
- (B) Hierarchical merger

- **B-DECIGO** can pin-down the story.
- Original observation.



# Observable Range

$30M_{\odot}$  BBH Merger : 100 Gpc ( $z > 10$ ) range  
with  $\text{SNR} \sim 8$  (optimal direction/polarization).







# B-DECIGO Sciences for CBC

---

- With its BBH observable range, in B-DECIGO  
Detection Rate will be  $\sim 4 \times 10^4 - 10^6$  events/yr .  
→ Possible to identify the origin of BBH :  
Pop-III, Pop-I/II, or Primordial BH.
- Range for BNS is  $\sim 2\text{Gpc}$  →  $\sim 100$  events/yr .
- With low-freq. GW observations, longer observation time is expected; in  $30M_{\odot}$  BBH merger case,  
the signal is at 0.1Hz in 15days before merger.  
→ Improved parameter estimation accuracy  
with larger cycle number ( $\sim 10^5$ ) :
  - \* Localization, Merger time → Alerts for GW-EM.
  - \* Mass, Distance, Spin → Origin and nature of BBH.



# Parameter Estimation Accuracy




T. Nakamura et al., Prog. Theor. Exp. Phys. 093E01 (2016)

# Target (3) : Foreground Understandings

In future DECIGO, unresolvable GWs by many binaries can be a foreground for primordial GW obs.

⇒ Gain understandings with >100 binaries.





# Technical Challenges

---


- Long-baseline Interferometry (Disp.  $< 2 \times 10^{-18} \text{ m/Hz}^{1/2}$ )
  - Optical configuration for IFO, and laser source.
  - 100km Fabry-Perot cavity (Large RoC, Distortion).
  - Initial attitude acquisition.
- Force Noise (Force noise  $< 1 \times 10^{-16} \text{ N/Hz}^{1/2}$ )
  - Gravity, EM force, Residual gas, thermal radiation, Cosmic ray, control noise, etc..
- Satellite control
  - Drag-free, Low-noise thruster, Signal processing.
- Satellite System Design
  - Orbital Design, Initial Mission sequence.
  - Resource distribution, Launcher, Cost estimation.

# JAXA Roadmap

内閣府・宇宙政策委員会・宇宙科学・探査部会 資料より (2013年9月19日).

## Ⅲ. 今後の宇宙科学・探査プロジェクトの推進方策

宇宙科学における宇宙理工学各分野の今後のプロジェクト実行の戦略に基づき、厳しいリソース制約の中、従来目指してきた大型化の実現よりも、中型以下の規模をメインストリームとし、中型(H2クラスで打ち上げを想定)、小型(イプシロンで打ち上げを想定)、および多様な小規模プロジェクトの3クラスのカテゴリーに分けて実施する。



2000年代前半までの  
典型的な科学衛星ミッション  
M-Vロケットによる打ち上げ

戦略的に実施する中型計画(300億程度)  
世界第一級の成果創出を目指し、各分野のフラッグ  
シップ的なミッションを日本がリーダーとして実施する。  
多様な形態の国際協力を前提。

公募型小型計画(100-150億規模)  
高頻度な成果創出を目指し、機動的かつ挑戦的に実施  
する小型ミッション。地球周回/深宇宙ミッションを機動的  
に実施。現行小型衛星計画から得られた経験等を活か  
し、衛星・探査機の高度化による軽量高機能化に取り組  
む。等価な規模の多様なプロジェクトも含む。


多様な小規模プロジェクト群(10億/年程度)  
海外ミッションへのジュニアパートナーとしての参加、海外  
も含めた衛星・小型ロケット・気球など飛翔機会への参  
加、小型飛翔機会の創出、ISSを利用した科学研究など、  
多様な機会を最大に活用し成果創出を最大化する。




# JAXA Roadmap

From file submitted to the government by ISAS/JAXA

(内閣府・宇宙政策委員会・宇宙科学・探査部会 2013年9月19日).




# DECIGO





# Updated Roadmap for DECIGO

Figure: S.Kawamura




# Space GW Antenna DECIGO

**DECIGO** (DECI-hertz interferometer Gravitational wave Observatory)

**Purpose: To Obtain Cosmological Knowledge.**


Direct observation of the origin of space-time and matter in Big-bang Universe.



## Interferometer Unit: Differential FP interferometer

Arm length:	1000 km
Finesse:	10
Mirror diameter:	1 m
Mirror mass:	100 kg
Laser power:	10 W
Laser wavelength:	532 nm

S/C: drag free  
3 interferometers






# Conceptual Design

## DECIGO

(DECI-hertz interferometer  
Gravitational wave Observatory)

Arm length:	1000 km
Finesse:	10
Mirror diameter:	1 m
Mirror mass:	100 kg
Laser power:	10 W
Laser wavelength :	532 nm

S/C: drag free  
3 interferometers



# Observation of the Early Universe



Background:  
original figure by  
NASA/WMAP Science Team




# Observation of GW from Inflation

BICEP2, (POLARBEAR, ...)

CMB B-mode polarization  
observation by micro-wave  
telescope.


DECIGO, (KAGRA, aLIGO, ...)

GWB observation by  
GW telescope.



# 'Window' for the Early Universe

DECIGO band is open window for **direct observation of the early universe.**



Kuroyanagi+, PRD (2009)


Pablo, PRD (2011)



# Probing the Early Universe by GW

---

- GWs will carry direct information on the early universe.
- Spectrum : Initial fluctuation + Evolution history



Depends on  $r$  (tensor-to-scalar ratio), which may be also pinned-down by CMB B-mode polarization observation.

Different age in different freq.  
Higher freq. → Earlier universe


- Reheating temperature
- Thermal history of the universe

...

# GW from Inflation

Energy density  $\propto$  Tensor-Scalar Ratio ( $r$ ).

Power spectrum : Evolution history of the Universe.



- Spectrum Power.  
→ Energy scale of inflation
- Cut-off freq.  
→ Energy scale of Reheating

Nakayama+,  
Journal of Cosmology  
and Astroparticle Physics  
06 (2008) 020.



---

# Summary

# Summary

---

- First direct detection of GW was achieved by LIGO 100 years after the theoretical prediction by A. Einstein by General Relativity.
- It opens the new field of 'Gravitational-wave astronomy'. We obtained a new prove to understand the universe.
- The field will be expanded by antennae with better sensitivity, and with different frequencies.
- Japanese KAGRA will improve the source parameter estimation accuracy. Best effort to join the network.
- B-DECIGO will provide fruitful sciences. Future DECIGO will be one of the dream of science; it will be able to observe the early universe directly.



# Related Parallel Sessions


---

GW1 (Mon): Ground-based GW detectors

GW4 (Tue): Mid.-frequency GW detection

GW5 (Thu): DECIGO

---



# End